

Historical Review of Nushagak River King Salmon Management

Prepared by

Tom Brookover

Anchorage, AK
tbrookak@gmail.com

Prepared for

Board of Fisheries Committee to examine the Nushagak-Mulchatna King Salmon
Management Plan

and

Bristol Bay Science and Research Institute

September 19, 2022

Table of Contents

INTRODUCTION	1
PRE-1987	2
KEY MANAGEMENT ISSUES	2
<i>Harvests and Exploitation rates.....</i>	<i>3</i>
<i>Gillnet mesh size and depth.....</i>	<i>4</i>
<i>Migratory behavior and timing.....</i>	<i>6</i>
<i>Inriver abundance and escapement assessment.....</i>	<i>7</i>
MANAGEMENT PROGRAM/TOOLS.....	8
RECOMMENDATIONS.....	9
<i>Habitat Protection.....</i>	<i>9</i>
<i>“Optimum” Escapement Goal.....</i>	<i>9</i>
<i>Estimation of Escapement</i>	<i>10</i>
<i>Achievement of Escapement</i>	<i>10</i>
SUMMARY, PRE-1987.....	11
DEVELOPMENT OF THE 1992 NUSHAGAK-MULCHATNA CHINOOK SALMON MANAGEMENT PLAN	12
PRE-PLAN, 1987-1991.....	12
DEVELOPMENT OF THE 1992 PLAN	13
POST-1992; PLAN CHANGES, FISHERY TRENDS, AND PLAN PERFORMANCE	15
PLAN MODIFICATIONS.....	15
COMMERCIAL FISHERY	16
<i>Regulation and Fishing Effort and Harvest.....</i>	<i>16</i>
SPORT FISHERY.....	22
<i>Regulations.....</i>	<i>22</i>
<i>Effort.....</i>	<i>23</i>
<i>Harvests</i>	<i>25</i>
SUBSISTENCE FISHERY	25
<i>Regulations, Effort, and Harvest</i>	<i>25</i>
PLAN PERFORMANCE.....	27
<i>Changes in Escapement Assessment Tool.....</i>	<i>28</i>
<i>Plan Objectives:.....</i>	<i>30</i>

MANAGEMENT CHALLENGES	32
ACKNOWLEDGEMENTS	32
LITERATURE CITED.....	34
APPENDIX A. 1992 VERSION, NUSHAGAK-MULCHATNA CHINOOK SALMON MANAGEMENT PLAN	38
APPENDIX B. 2019 VERSION, NUSHAGAK-MULCHATNA KING SALMON MANAGEMENT PLAN..	39
APPENDIX C. TABLES.	41

Introduction

In 1992, the Alaska Board of Fisheries (Board) adopted the Nushagak-Mulchatna King Salmon Management Plan (Plan) to guide management of the subsistence, commercial and sport fisheries that harvest this important stock. The Nushagak River fisheries that harvest Chinook (king) salmon have been managed under the direction of the Plan since then. However, restrictions to the sport fishery due to low early season inriver passage of king salmon combined with sometimes intense fishing for sockeye in the Nushagak District in the mid-2010's led to calls to pair restrictions in the commercial and sport fishery in 2018. Proposals 41 and 42, submitted for deliberation at the November 2018 Bristol Bay Board meeting, both sought to restrict time in the commercial fishery when the sport fishery is restricted in season by emergency order.

In response to the proposals, the Board established a committee at the 2018 meeting to develop a comprehensive solution to the Plan and charged the committee with reporting back to the Board. The Bristol Bay Science and Research Institute (BBSRI) also committed to supporting the committee's work through a stakeholder-led technical analysis of options the committee was expected to consider. Possible committee products included regulatory proposals and/or other non-regulatory recommendations.

An early (October 14, 2019) draft version of this report was developed to summarize management of Nushagak River king salmon for the committee's benefit. The history of the fishery through the mid-1980s was well documented in a comprehensive, albeit dated, report (Nelson, 1987). The 2019 draft of this report provided an updated comprehensive historical overview summarizing Nelson's report as a basis, then describing the evolution of the fisheries that followed.

The purpose of the 2019 draft was to provide committee members with key information, help create a better understanding, and provide a basis for future recommendations concerning management of the Nushagak River king salmon fisheries. The draft was intended as a "living" document and was expected to evolve with input from committee members and others and as new fishery information came available.

The committee met initially October 21, 2019, in Anchorage to get underway and discuss preliminary analysis of the fishery's history, including information presented in the draft report, and technical challenges associated with the monitoring and management of the fishery. Break-out groups met in December 2019 and February 2020. At the Upper Cook Inlet meeting in February 2020, the Board disbanded the formal committee but encouraged stakeholders on the committee to continue to work together in preparation for the next in-cycle Bristol Bay meeting. Since then, the committee met on numerous occasions toward developing comprehensive recommendations to improve the Plan and stock assessment programs in preparation for the Bristol Bay Board meeting scheduled for November 2022. BBSRI facilitated the meetings and provided technical analysis and support. The committee

process and outcomes are to be discussed in depth in a separate report and are therefore not discussed in this one.

In this report, historical king salmon management in the Nushagak District is portioned into three eras:

- 1884-1986 (recap of Nelson (1987))
- 1987-1992 (development of the Plan)
- 1992 through 2021 (the Plan years)

This report includes fishery data for the years that followed the early draft (2019, 2020 and 2021). Discussion of fishery trends have been adjusted accordingly. Comments received from committee members and staff from the Alaska Department of Fish and Game (ADF&G) have also been incorporated. The report is intended to be made available with other work products, including a separate report on the committee process and a proposal to the Board detailing changes to the Plan, to the public prior to the 2022 Board meeting. Like the 2019 draft, its purpose is to improve understanding of the Nushagak River king salmon fisheries and their management and provide a basis for committee recommendations.

Pre-1987

The history of the Nushagak king salmon fisheries from the inception of the commercial fishery in Nushagak Bay in 1884 through the mid-1980s was well documented in a comprehensive report (Nelson, 1987). Mike Nelson worked as the Area Biologist for the ADF&G in Dillingham and oversaw management of the Nushagak commercial and subsistence fisheries from shortly after statehood until his retirement in 1987. The purpose of the report was to assist in creating a better understanding of the king salmon management program and provide a basis for future recommendations regarding fishing regulations. Nelson (1987) helped set the stage for the development of the Nushagak-Mulchatna King Salmon Management Plan in 1991.

This section summarizes Nelson's findings. By the time the report was published, the commercial fishery had "traditionally extracted a heavy toll from the total run, while freshwater sport fishing interests (were) growing rapidly." There was a growing concern that spawning escapements may be jeopardized, and that the natural productivity could not be maintained. As greater fishing pressure was exerted on the stock, the fisheries were subjected to progressively more stringent regulations. Under this background, Nelson foresaw a clear need for "a careful, quantitative appraisal of the fishery impacts and of regulatory options" to maintain or increase productivity and address hardships among the various participants.

Key Management Issues

Nelson (1987) clearly recognized the value of Nushagak River king salmon to the area's commercial, subsistence and sport fisheries, as well as the challenges presented by then-

apparent very high exploitation rates and fishery practices. These included the potential for friction among the fisheries in the face of increasing demand as well as conservation-related concerns for the quantity and quality of escapement and resultant impacts to productivity of the stock. Several salient points discussed in the report included:

- exploitation rates had exceeded 95% of the early run component and were expected to remain high without further restrictions,
- gill net mesh size and depth directly influenced exploitation rates and quantity and quality of escapement,
- fish holding within and above the district created difficulties in obtaining escapement throughout the run, and
- methods to assess inriver abundance/spawning escapement were under development

Each of these points are discussed in more detail in the following sections.

Harvests and Exploitation rates

The commercial fishery for salmon in Bristol Bay began in 1884. Sockeye salmon were, and remain, the targeted species and main emphasis for the Bristol Bay and Nushagak fishery. However, the commercial harvest of king salmon in the Nushagak District advanced rapidly once development began. After sustained commercial utilization (1955-1971), catches declined (1972-1975) but recovered, and then reached a historical peak over the decade 1976-1986. Recovering salmon markets and advances in gear effectiveness at catching king salmon were primary factors driving the renewed commercial interest in early season fishing effort. However, peak production of king salmon in the early 1980s resulted in a surge of interest and record harvests in the commercial fishery. Nelson (1987) chronicles the trends in commercial harvest from the fishery inception through 1986; annual harvests ranged from 1,635 (1935) to 195,287 (1982) fish with the three largest harvests occurring in 1979, 1981 and 1982. By 1987, the Nushagak watershed produced the state's second largest stock-specific commercial king salmon fishery, nearly matching those of the Yukon River.

He similarly discussed trends in the subsistence and sport fisheries. While subsistence use of salmon dated back beyond the availability of written literature, little data on harvest was available prior to 1963 when a permit system was initiated. Subsistence harvests in the Nushagak District normally ranged between 50 and 80 thousand salmon and had been increasing due to increased effort from local population increases and annual influxes from non-watershed participants, and better harvest reporting. As king salmon are the first species to arrive in the spring, they received considerable interest and fishing pressure. From 1963 through 1986, subsistence harvests averaged 7,200 and ranged from 2,900 (1964) to 12,600 (1986) king salmon. Effort and harvest of king salmon had increased since 1970 and, like the commercial fishery, the subsistence fishery accounted for its largest harvests in the early 1980s.

Development of sport fisheries in Bristol Bay had occurred more recently relative to commercial and subsistence fisheries. Nelson cited Paddock (1964) describing the first significant instance of king salmon sport use on the Nushagak River taking place at Portage Creek in 1963. Since then, sport fishing had become more popular in Bristol Bay, and the peak production of king salmon in the early 1980s contributed to the growing fishery on the Nushagak River, with increasing effort and harvest. Sport harvests were estimated from 1977 to 1986. The largest sport harvest occurred in 1984 (2,382 fish).

Using available catch and escapement data from 1966 through 1986, Nelson (1987) estimated the average Nushagak king salmon total run at over 176,000. He noted an improvement in the adult production trend whereby then-recent runs (1978-1986) averaged 246,000 fish, nearly twice the size of runs averaged from 1966-1977 (125,000 fish). Over the entire period, exploitation rates averaged 54 percent and ranged from 29 (1975) to 72 percent (1969).

Exploitation on the early component of the king salmon run appeared to be of specific concern; then-recent commercial and subsistence exploitation rates had exceeded 95% for this component. Traditionally, the commercial fishery commenced in late May to early June. Approximately 85% of the annual harvest was taken in the month of June and the mid-point was June 18. Nelson (1987) describes a bimodal pattern of harvests taken 1973-1986, with the first peak occurring June 7-14 and the second, June 23-26. He ascribes the bimodal pattern to the established fishing schedule of 5 days per week prior June 16, when the fishery was closed unless opened for fishing by emergency order and notes that, as more pressure was exerted early in the run, fishery managers applied additional time and area closures. The effect of those actions became apparent in 1981, when high catch rates shifted from early in the season to later.

Gillnet mesh size and depth

Gillnets were (and remain) the only fishing gear allowed in the commercial fishery and were the only gear used in the subsistence fishery. Drift gill net gear accounted for most of the total catch. As a result, and because of the characteristics of the gear related to fish size regardless of species, Nelson (1987) focused considerable discussion on the impacts gillnet mesh size and depth have on king salmon.

By 1987, basic data on age, weight and length had been collected from the Nushagak king salmon harvests and spawning escapement. According to Nelson (1987), a statistically adequate number of samples had been collected each year from the commercial fishery beginning 1966, and from subsistence harvests and spawning escapements beginning 1982. Based on analysis of the samples collected, Nelson (1987) described some of the biological characteristics of Nushagak king salmon as follows:

- Age class composition of the run varies from year to year; however, most king salmon (80 percent) return as 5- and 6-year-old fish and over 96 percent return as age 4 through 7.

- Age class differences between males and females is striking; age 4 and 5 fish are predominantly males and in contrast, age 6 and 7 fish are predominately females.
- Based on data from the commercial fishery, there is considerable overlap of lengths between age classes. Females are generally longer than males of the same age class through age 6.
- Mean weight of females tends to be greater for a given age class compared to males.
- Age at sexual maturity varies between males and females.
- A weighted average (1982-1984) of catch and escapement indicated a higher proportion of males (53 percent) in the total runs.
- Based on fecundity data collected from the 1966 and 1968 Nushagak District commercial catches (n=69), number per female averaged over 10,000 eggs. Nushagak River king salmon appeared to have some of the highest fecundity rates found in the species throughout the Pacific Coast.

At that time, the Nushagak gill net fishery showed considerable selectivity by age, size, and sex. Historically, large mesh nets were used to target king salmon while smaller mesh nets were used to target sockeye salmon. Gillnet specification varied from year to year but by the mid- 1970s, 8 to 8 ½ inch mesh was commonly used to target king salmon (early in the season), while sockeye salmon were targeted using 5 1/8 to 5 ½ inch mesh gillnets (later in the season). Smaller mesh nets (5 3/8 inch) tended to selectively capture smaller king salmon which are primarily males, while larger mesh nets (8¼ to 8½ inch) tended to select for larger salmon which are primarily females. Thus, early season (large) mesh accounted for a heavy preponderance of large females in the catch, while smaller mesh sockeye gear accounted for a higher proportion of younger age males. Some important additional points regarding mesh selectivity made by Nelson (1987) follow:

- The commercial fishery showed an overall higher percent of males which Nelson attributed to a relatively greater abundance of early maturing, smaller age 4 and 5 males.
- Mesh selectivity affected the age and sex composition of the escapement.
- A weighted average (1982-1984) of catch and escapement indicated a higher proportion of males in the catch and a higher proportion of females in the escapement.
- Since large mesh gill nets tend to harvest larger female fish, mesh selectivity affected the average fecundity of the female spawning population. King salmon harvested with large mesh, i.e., 8 ½ inch, nets vs small mesh, i.e., 6 ½ inch, nets resulted in a two-fold difference in egg deposition on the spawning grounds.
- Large mesh gill nets were restricted for the first time in 1985 and 1986 to reduce catch rates and were felt to be effective in allowing additional large king salmon into the river to spawn.

While mesh size restrictions were historically implemented to manage sockeye salmon harvest, then-recent use of inseason restrictions on the use of large mesh showed promise in reducing exploitation of large fecund females.

Nelson stated that gillnet (mesh) depth was of equal importance to mesh size with respect to catch rates for king salmon. King salmon appear to follow deeper water channels in the generally shallow waters of the Nushagak District, where deeper nets are more effective.

Gillnet length and mesh size varied during the early years of the fishery until 1923 when the U.S. Bureau of Fisheries restricted both. At the time of the report, little information existed on the depth of king salmon nets in existing literature, and the depth used appeared to closely follow a 28-mesh restriction enacted in 1925 for sockeye salmon nets.

As interest in king salmon increased in the 1940's, some Nushagak fishermen began to experiment with deeper nets. Reports from fisherman indicated higher success rates with deeper nets through the mid-1950s and, as fishermen became more effective with deeper nets, interest and participation in the fishery accelerated.

By 1957, Federal fishery managers recognized that the increase in fishing effort required additional closed time for king salmon conservation purposes. In 1958, weekly fishing time (prior to June 22) was reduced by 36 hours and nets were limited to 28 meshes in depth. Nelson cited an experienced fisherman attesting to effectiveness of the depth restriction in reducing the increased exploitation on and stated that the depth restriction is an essential component of the regulatory management program for the species.

Migratory behavior and timing

Nelson made the point that, considering the rapid growth and "gross mismanagement" of the early Bristol Bay sockeye salmon fishery, Nushagak king salmon were fortunate in that the run arrived before the sockeye fishery began in earnest. Thus, the advanced (earlier) run timing of the species, along with the relatively low commercial interest in its smaller run, helped the stock survive the development of the sockeye fishery.

Fishery managers began to use this difference in timing to manage for conservation of king salmon in 1958. When weekly fishing time was reduced and net depth was restricted that year, the restrictions were applied prior to June 22 when king salmon were the primary species present. As fishermen became more effective at targeting king salmon and effort targeting the species increased, fishing time prior to June 16 was further reduced. For the 1987 season, ADF&G planned to prohibit fishing prior to June 1 and replace the 5-day fishing schedule then in place prior to June 16 with a 3-day schedule. At the time, fishing beginning June 16 was closed unless and until opened by emergency order. Future action, including replacing the fishing schedule prior to June 16 with emergency order management, would be considered depending on the success of the 1987 measures.

While the earlier run timing relative to sockeye salmon contributed to king salmon sustainability and provided a means to manage the species separately for conservation, other migration tendencies posed management challenges. King salmon often mill and hold within the district, are believed by many fishermen to hold deep during calm weather and therefore unavailable to the fishery and appear to move upriver and become available to the fishery under the influence of strong winds. For these reasons, the effectiveness of early

season closures on reducing harvest rates was limited at times; early season closures coincided with a noticeable shift in high catch rates from early to later in the season in the early 1980s.

Run timing data was collected from four sources: commercial, subsistence and sport harvests, and sonar-based enumeration. Over half (55 percent) of the commercial harvest was accumulated by June 16-20. Subsistence harvest in the Dillingham area peaked between June 20-30 (later upriver). Sport catches inriver peaked between June 26 and July 6. And available sonar data indicated 50% of the inriver run had passed the sonar site July 1-2. Nelson acknowledged the commercial fishery can influence the migration timing of the inriver run but pointed out that the data collectively indicated that most king salmon migrate into the lower river during late June to early July.

Inriver abundance and escapement assessment

Management of salmon fisheries in Alaska is based primarily on achieving escapement levels that support sustainable harvests. As Nelson stated: “the criterion of escapement has been the primary factor in determining fishing regulations in Alaska, from the passage of the White Act in 1924 to the present time.” Yet, the magnitude (and quality) of spawning escapements has not always been estimated. Escapement data for king salmon is relatively difficult to collect because spawning is generally concentrated in mainstem reaches of larger, turbid river systems.

Aerial surveys to locate king salmon spawning areas and assess spawning magnitude in the Nushagak River began in 1956 and continued through publication of the report (and beyond). One of the objectives of the aerial survey assessments was to develop methods to use aerial survey counts to estimate total escapement.

In 1979, a side scanning sonar project to enumerate adult sockeye salmon was initiated on the lower Nushagak River near Portage Creek. Nelson acknowledged the potential of the sonar project to estimate king salmon escapement but continued aerial surveys during the subsequent years due to operational difficulties and sampling problems experienced by the sonar project. Some of the initial challenges of using sonar to estimate passage included exceeding the density threshold of the Bendix units, limited sonar range/coverage of the migratory pathway of the larger king salmon, and difficulties in apportioning sonar targets to specific species among the sockeye, chum, and king salmon that comigrate past Portage Creek.

Annual monitoring of daily subsistence catches at Lewis Point on the lower Nushagak River was initiated in 1980 to provide daily estimates of king salmon escapement in advance of estimates provided by the sonar project. Unlike aerial survey assessments conducted on the spawning grounds, both the sonar and Lewis Point catch monitoring projects provided the added benefit of inseason “real-time” data on inriver abundance in the Nushagak River. However, problems with the Lewis Point project also kept the emphasis on the aerial survey program as the primary means to estimate spawning escapement.

Visual counts of salmon passing by points on the shoreline were conducted from counting towers beginning in 1953 to estimate sockeye escapement. Incidental tower counts were also collected routinely for king salmon. Counting periods, designed to capture the duration of the sockeye run, did not cover the duration of king salmon run and counts were of limited use as a result. One weir project – 1968 Stuyahok River weir - had been implemented in Bristol Bay to enumerate king salmon.

Beginning in 1966, an expanded ‘comprehensive’ aerial survey program was used to expand counts of king salmon to total inriver spawning abundance. Expansion factors and methodology varied by year and had not been rigorously evaluated until 1982 after an extensive series of escapement data had been collected from numerous spawning streams within the Nushagak drainage. In that evaluation, selected portions of the Nushagak and Mulchatna main stems, for which counts had been collected for eight years, were correlated with total counts for years when they were available. The correlation, in turn, was then used to estimate total escapement in the Nushagak drainage. Resulting escapement estimates from 1966-1986 averaged 82,000 and ranged from 25,000 (1972) to 162,000 (1983).

Management Program/Tools

Unlike the Bristol Bay sockeye salmon fishery, the Nushagak king salmon fishery received little directed effort at research and management until the 1950s. In the 1960s the management strategy was to limit harvest to a range of 60,000 to 80,000 fish with exceptions. As pressure on king salmon increased in the 1970s, the need for more robust escapement data collection also increased. And as the sport fishery grew so did the need for information on sport fishing use. In addition to funding and staffing the Dillingham area office with biologists and technicians assigned to commercial and sport fish management and research in the Nushagak District, ADF&G conducted a suite of programs aimed at king salmon at the time the report was written:

- Commercial and subsistence harvest monitoring – daily contact with processors enabled commercial catch estimates and harvest rates. Project objectives included inseason estimates of catch and fishing effort for king salmon by period, and inseason catch per unit effort.
- Commercial catch sampling – king salmon from commercial harvests were measured for weight and length, sex determined, and scale removed for age determination. Project objectives were to provide age, weigh, length, and sex data for commercially harvested king salmon.
- Sport fishery harvest monitoring
 - Creel surveys in the lower Nushagak River – anglers were interviewed inseason to collect catch and harvest data, and sample harvested fish. Project objectives included estimates of angling effort, catch and harvest rates, and collection of biological and demographic data.
 - Statewide Harvest Survey – postal surveys were mailed annually to anglers that fished in Alaska to collect effort and harvest data. Results provide harvest estimates for the Nushagak king salmon sport fishery.

- District test fishing – Fishing with gillnets took place within the Nushagak District to capture salmon. The primary objective was to monitor magnitude and entry pattern of sockeye salmon in the district. A secondary objective was to provide indications of when king salmon were present, holding, and moving upriver of the district.
- Lewis Point subsistence/test fishery – Lewis Point subsistence catches were monitored and sampled. Objectives were to estimate escapement into the river using subsistence catches, and sample catches for age, sex, and length data.
- Post-season aerial surveys – comprehensive surveys were flown to count spawning king salmon. Primary objectives were to provide estimates of drainage-wide escapement and spawning distribution.
- Portage Creek Sonar – obtain daily salmon passage rates from two Bendix side-scanning sonar units in the lower river near Portage Creek, sample salmon for age, sex, and length data, and adjust sonar counts by species. Project objective was to estimate inseason escapement of salmon by species.

At the time Nelson (1987) was published, data collected from these projects were used for king salmon inseason fishery management, post-season management assessment, and beginning in 1984, pre-season forecasts of projected run size.

Recommendations

Nelson (1987) identified four categories of needs that should be addressed: habitat protection, optimum escapement objectives, methods to accurately estimate escapement, and methods to achieve escapement objectives.

Habitat Protection

Nelson described the protection of freshwater spawning and rearing habitat a priority requirement to sustained and increased king salmon production. Three habitat objectives were identified as referenced from the 1986 Comprehensive Salmon Plan:

- Maintain present quantity and quality of salmon habitat
- Enforce state water quality and anadromous stream protection regulations, and
- Develop land use plans for public lands adjoining salmon waters

“Optimum” Escapement Goal

Although provisional escapement objectives were in place, Nelson indicated a final goal should be developed and suggested delaying its development until after the 1990 run, when returns from the large escapements in 1981-1983 would be complete.

- Develop an optimum¹ escapement goal (after 1990 run)

¹Nelson used the term *optimum* escapement goal like the way we currently use biological escapement goal (BEG) based on expected maximum sustainable yield (MSY). He did not use it to mean the same thing as today’s Optimum Escapement Goal (OEG) in the State’s escapement goal

- Continue to collect age, sex, length, and weight data needed for escapement goal development and run forecasting
- Conduct a mesh size study to determine the effects of mesh size on reproductive potential, and assess the use of regulatory mesh size restrictions as a king salmon management tool
- Conduct a tagging study to assess movement and holding patterns in the fishery, district, and lower river.

Estimation of Escapement

Nelson envisioned substantial benefits to providing more accurate and timely information with which to estimate inseason escapement rates. Primary benefits included allowing for additional harvest during strong runs while providing additional protection to smaller runs.

- Improved subsistence monitoring, i.e., test fish project at Kanakanak Beach, to provide daily catch estimates and possibly additional data
- Continued development of the Portage Creek sonar to provide inseason and total estimates of escapement. Species apportionment was the primary challenge to reaching this objective. Successful development would allow the termination of the aerial survey program.

Achievement of Escapement

This goal was aimed at providing managers with effective methods to control fishing pressure and achieve escapement goals. It was predicated on defining optimum escapement objectives and developing methods to accurately estimate inseason escapement rates.

- Conduct the commercial fishery entirely under day-to-day (emergency order) management if planned regulatory changes in 1987 were not effective in reducing the exploitation rate to achieve better distribution of escapement through time.
- Restrict large mesh gill net gear to reduce catch rates

Finally, Nelson noted positive attributes of the Nushagak king salmon stocks compared to others in Alaska: the stock is generally in good condition; is concentrated in a large river system that can be managed independently; the fisheries on the stock are conducted in a terminal area where allocation considerations are modest and, king salmon are somewhat separated from other species by timing differences in most years. Ultimately, he noted: *“the success of management will depend on the effectiveness of stock assessment capabilities and maintenance of a management strategy that is responsive to stock abundance, while retaining an element of conservatism in response to uncertainty about stock productivity.”*

policy, which is set by the Board of Fisheries and takes into account biological and socio-economic factors to set the escapement goal target.

Summary, Pre-1987

The period from the early 1950s through 1986 was formative in the development of the Nushagak fisheries and their management. The period experienced a growing interest in Nushagak River king salmon, and peak production of king salmon enjoyed in the early 1980s resulted in a surge of interest and record harvests in the commercial fishery, and development of a growing sport fishery. Together, these dynamics presented concerns for adequate spawning escapement and potential for user conflicts.

Fishery managers responded to the increase in interest by enacting fishery restrictions to ensure enough king salmon for spawning escapement. In 1958, Federal fishery managers had restricted weekly commercial fishing time and gillnet depth to boost the escapement. Subsequent restrictions to fishing time, area and gear were implemented by state managers through the mid-1980s. In 1985 and 1986, large mesh gill nets were prohibited by emergency order. Plans for 1987 called for reducing area in the outer district, prohibiting fishing before June 1, and reducing the weekly fishing schedule prior to June 16 from five to three days.

Fishery managers also responded to the increased interest in the fishery by adding stock assessment programs to ensure conservation of Nushagak king salmon. Aerial surveys to document escapement began in 1956. In the 1960s, State managers expanded the aerial survey program to additional systems within the drainage and implemented a subsistence permit system in part to provide better accounting of subsistence fishing activity. In 1979, the side-scanning sonar project at Portage Creek was implemented to enumerate sockeye salmon with an interest in using that system to index or enumerate king salmon. In the 1980s, creel surveys were initiated to estimate sport fishing effort and harvest.

Improved stock assessment allowed for additional tools to use in managing the Nushagak king salmon fishery. By 1987, fishery managers had compiled a time series of estimated harvests for each fishery component and escapement, which allowed for annual estimates of total run size. Age composition estimates obtained for each component allowed for the development of brood tables, which in turn provided information needed to develop a biological escapement goal and, beginning in 1984, an annual pre-season forecast of the run.

Despite the advances in stock assessment and increasingly conservative management of the fisheries, conservation issues remained to be addressed as of 1987. A formal escapement goal had yet to be developed. Accurate and timely (daily) inseason escapement estimates, needed to take advantage of harvestable surplus of large runs and conserve small runs, required continued research and development of the sonar program at Portage Creek. Species apportionment of fish counted by sonar continued as a major obstacle to inseason assessment. Finally, managers recognized that additional management measures may be needed should the restrictions envisioned for 1987 not be effective enough to control fishing pressure and achieve escapement objectives.

Development of the 1992 Nushagak-Mulchatna Chinook Salmon Management Plan

Pre-Plan, 1987-1991

While the period spanning the 1950s to the mid-1980s was formative in the development of the fisheries and their management, the following several years cemented the need for a structured management plan. A weak king salmon run in 1986, coupled with a poor forecast for the 1987 run, indicated that the large runs experienced in the late 1970s and early 1980s were coming to an end (Minard et al., 1992). Indeed, runs observed from 1987 through 1990 (range 86 to 146 thousand) declined from the very large runs observed from 1978 to 1983 (range 218 to 356 thousand) to a level generally considered as 'depressed'.

By 1991, it had become evident that the large runs experienced in the early 1980s had produced poorly; spawning escapements from brood years 1981-1985 had produced only as many fish as had spawned in those years, or fewer. After a comprehensive review of production data, Minard et al. (1992) stated that the decrease in production at higher escapement levels was the most notable trend in the spawner-return data. Normally, this would indicate density-dependent factors in the freshwater environment. However, in this case where large escapements all occurred sequentially among brood years 1981-1985, it is difficult to determine whether the decrease in production was caused by the high levels of escapement or by other factors that may have occurred during the life cycle of salmon produced in those years (e.g., changes in ocean carrying capacity, high seas fisheries interceptions, freshwater habitat degradation, competition with other species in the fresh and/or marine environment).

The return to more typical (or depressed) run sizes in the mid-1980s prompted managers to implement additional conservation measures. These included emergency order management of the commercial fishery that Nelson had suggested, which ultimately led to closure of the directed commercial fishery. The 1987 commercial fishery opened normally but was closed by EO after approximately 5,000 king salmon were caught with little indication of fish movement into the river. The commercial fishery was similarly closed by EO each of the three subsequent years, prompted by low pre-season forecasts and a likelihood of large incidental harvests of king salmon in the sockeye fishery. An improved forecast in 1991 and indications of escapement more than the goal prompted a commercial period June 24, 1991. However, a boycott by commercial harvesters over salmon prices kept fishing effort low.

During this period, the Board of Fisheries implemented several conservation measures affecting the commercial and sport fisheries.

- Prior to the 1988 season: the outer king salmon boundary was eliminated by regulation; the commercial district was redefined to include only the sockeye salmon boundary as the southern-most district boundary line. This effectively reduced potential fishing area for king salmon.

- the regulatory commercial fishing season was reduced from May 1 to June 1.
- sport fishing bag limits in the Nushagak drainage were reduced from 5 king salmon per day and in possession, of which only 2 may be over 28 inches, to 3 king salmon per day and in possession, of which only 2 may be over 28 inches.
- The following year (1989), the Board abolished the minimum mesh size requirement of 6 ¾ inch mesh in place in the commercial fishery prior to June 16.
- In 1990, the Board closed the Nushagak River drainage upstream from its confluence with the Iowithla River, including the Iowithla River, to the taking of king salmon from July 25 through December 31.

The poor runs experienced during this period underscored the need for a revised escapement goal as recommended by Nelson. Other dynamics further heightened the need. The provisional escapement goal was not attained in 1986, 1988, and 1990. Additionally, commercial salmon fishery managers in Bristol Bay had traditionally accounted for returns as either commercial catch or escapement, the notion being inriver harvests were so small that their impact on inriver abundance was insignificant. With growth in the subsistence and sport fisheries, and ADF&G's mandate to manage for sustained yield, inriver harvests had to be explicitly accounted for in the escapement goal. This meant that the provisional 'escapement' goal of 75,000 was an inriver goal, and by managing for 75,000 fish at the Portage Creek sonar, the goal of attaining a spawning magnitude of 75,000 king salmon would not be realized.

Nelson (1987) described concerns with the heavy toll extracted by the commercial fishery and the growing sport fishery, and identified the need for improved escapement monitoring, a formal escapement goal, and additional management measures for the Nushagak king salmon fisheries in 1987. The poor performance of the large escapements during the early 1980s, the increasingly severe restrictions in the late 1980s resulting from the depressed runs, and the state of the provisional escapement goal all heightened concerns over conservation and exacerbated user conflicts that had begun to develop prior to 1987. During this period, they were raised to a level that received the attention of fishery participants, managers, and regulators alike, and turned the heat up on the need to develop and implement a formal management plan. Because such a plan would affect allocation among users, it had to be developed via the Board of Fisheries process to be effective.

Development of the 1992 Plan

Prior to the 1992 Bristol Bay Board meeting and under correspondence from the Board, the Nushagak Advisory Committee (NAC) submitted Proposal 157, and ADF&G submitted Proposal 158 to develop a management plan for Nushagak River king salmon. Both proposals expressed concern over poor recent runs and poor production trend and a need to provide ADF&G with management direction. The NAC proposal specified high seas bycatch and interception as a concern (but recognized that the issue was outside of the scope of the Board of Fisheries), and referenced habitat degradation and inriver harvest as possible factors influencing low return rates. The ADF&G proposal recognized the need to change the escapement goal to better account for biological needs and upriver harvests.

In support of the planning efforts, ADF&G conducted a review of the then-present escapement goal (Minard et al. 1992). Estimates for number and age of king salmon harvested in each fishery and for spawning escapement were available with limitations, and significant assumptions were made regarding the applicability of the data. Estimates of “biological escapement requirement” (BER), what we would call a Biological Escapement Goal (BEG) today, were derived using multiple methods, and ranged from 50,000 (early-years Ricker model) to 65,000 (all-years Ricker model) king salmon spawners. ADF&G recommended a BER at the upper end of this range to be conservative because of uncertainty in the brood tables and the uncertainty over the cause of the poor returns from the 1980-1985 runs.

Both the NAC and ADF&G proposed developing a plan that would distinguish inriver harvests from the BER, include management guidelines developed by the Board to share the burden of conservation among fisheries and provide staff with management direction, and achieve the BER. The NAC proposal prescribed specific management measures for each fishery under various projected escapement levels. Both proposals recognized that: “without a well described management plan, continued exploitation by the user groups on an apparently declining stock could have a long-term negative affect on this important stock.”

Prior to the January 1992 Board meeting, ADF&G and the NAC worked together on further developing a plan. By December 1991 the committee with ADF&G’s assistance had developed a draft (December 18, 1991) that contained much of the structure and content ultimately adopted by the Board in January 1992. The December 1991 draft included a BER of 65,000 spawners established by ADF&G during the then-recent escapement goal review. It included an inriver goal of 75,000 king salmon to provide for the BER and subsistence and sport harvest occurring upstream of the sonar. And it included management measures for the fisheries under three tiers based directly on projected inriver abundance estimates at the sonar.

Using the NAC draft plan as a template, the Board of Fisheries deliberated over the course of two days and approved the Nushagak-Mulchatna King Salmon Management Plan January 8, 1992 (Appendix A). The Plan directed ADF&G to manage the commercial fishery to achieve an inriver goal of 75,000 king salmon upstream from the Portage sonar site. The inriver goal provided for a BER of 65,000 and harvests above the sonar in the subsistence and recreational fishery. The Plan also set a cap on the recreational harvest not to exceed 5,000 king salmon.

The Plan was structured under three tiers and associated triggers tied to projected inriver run levels, much as it is remains today.

- At projected runs less than 40,000 king salmon, the sport and directed commercial fisheries were to be closed, the commercial fishery for sockeye was to remain closed until 10% of the Wood River escapement goal is projected, and the subsistence fishery was to be restricted by time or area.

- At inriver runs projected between 40,000 and 75,000, the directed commercial fishery for king salmon was to be closed and gillnets with greater than 5 ½ inch mesh were to be prohibited. At inriver runs projected between 40,000 and 65,000, sport fishing was to be restricted.
- At projections above 75,000 the Plan called for no restrictions on the commercial or subsistence fishery. However, at projections from 75,000 to 95,000 the sport fishery was to be managed such that harvests did not exceed 6,000 king salmon.

The third tier, in which inriver runs are projected to exceed the inriver goal, received considerable attention at the board meeting. The ‘cap’ on the sport fishery was one of the more controversial elements of the Plan. Some considered capping the sport harvest when harvestable surplus was available as consistent with the purpose of harvesting king salmon in the fisheries that historically harvest them. Others argued that capping sport harvest at or above optimum levels of yield was inconsistent with the sustained yield principle, particularly after other fisheries are afforded harvest under the same scenario.

Post-1992; Plan Changes, Fishery Trends, and Plan Performance

Thirty years have now passed since the Board adopted the original Plan. Over time, changes have occurred in the Nushagak king salmon commercial, subsistence and sport fisheries and the Plan. This section is intended to highlight some of the key dynamics in the fisheries governed by the Plan since 1992 and characterize how the Plan has performed relative to its stated objectives over time.

Plan Modifications

The Plan has been modified seven times by the Board of Fisheries (Table 1). Its purpose and structure, with management actions directly based on inriver run projections to the sonar, has remained very similar to the original version.

Management trigger levels (inriver projection levels of 40,000, 65,000, 75,000 and 95,000 king salmon) have changed twice. The first, in 1997, was specific and effectively reduced the range in which sport fishery restrictions were to be issued from 40,000-65,000 to 40,000-55,000. The 55,000-fish trigger was adopted partly based on analysis that showed little difference in expected productivity between the two levels. In addition, the 65,000-fish trigger had become disruptive to the sport fishery by precipitating frequent inseason restrictions prior to 1997.

The second, in 2012, changed the inriver and escapement goals and all management triggers contained in the Plan. The Board made these changes as requested in a proposal submitted by ADF&G to reflect a transition/conversion from Bendix to DIDSON sonar, because DIDSON accounted for a higher proportion of the king salmon that migrate up the Nushagak River. The biological escapement goal was changed from 65,000 to a range of 55,000-120,000 king salmon, the inriver goal was revised from 65,000 to 95,000 king salmon, and the various management triggers were changed as well.

Other changes to the Plan are discussed under the relevant fisheries below. The current Plan can be found in Appendix B.

Commercial Fishery

Regulation and Fishing Effort and Harvest

Directed commercial fishing for king salmon resumed under the Plan in 1992 (Table 2). Decisions to open the directed fishery and set the opening durations were based largely on the pre-season forecast and inseason indicators of run strength, including commercial harvest performance, subsistence harvest rates, an inriver passage rates estimated at the Portage Creek sonar (Brookover et al., 1997; Morstad et al., 2010).

The approach to scheduling directed openings varied from 1992 to present. Initially, the number and duration of openings were limited. Openings were generally scheduled to follow inriver pulses of fish evidenced by spikes in subsistence catch rates and other indicators (Brookover et al., 1997). This ensured fish migrate inriver prior to exposure to the commercial fishery. From 1994 to 1996, the directed fishery was managed more aggressively to harvest available surplus by scheduling more openings during lulls in fish passage. However, due to escapement quality problems observed in 1995 and 1996, commercial fishing periods in 1997 were scheduled directly after pulses of fish were observed moving into the river again, to reduce selectivity for large fish. The Board subsequently modified the Plan directing ADF&G to schedule openings to provide pulses of fish into the river that haven't been subject to harvest with commercial gear. From 2003 through 2009, the management strategy included openings earlier in June, with more space between openings, when a surplus appeared to be available (Fair et al., 2004; Westing et al., 2005, Morstad et al., 2010). Opening early in June during the first third of the run was intended to allow for lower levels of harvest over a larger portion of the run, still provide for fish movement past the district, and provide improved market quality and value to fishermen but carried the potential of overharvesting the early part of the run. Beginning in 2010, stakeholder meetings were used to help establish directed fishery schedules prior to the season (Salomone et al., 2011).

From 1992 through 2010, the directed commercial fishery was opened every year except two (2000 and 2001; Figure 1). Commercial fishing opportunity, based on the number of openings and total fishing time, was highest during 1994, 1995, 1998, and 2005-2007. During the 1990s, 200 or more drift boats participated based on boat counts conducted during the open fishing periods, with the largest boat counts recorded in 1994 and 1995. As an indication of the popularity of the directed fishery, the peak daily commercial drift permit registration for the 1994 and 1995 seasons occurred on dates during the directed fishery; in all other years the peak daily registration for the season occurred during the sockeye salmon fishery (Table 3). Number of drift deliveries peaked in 2005 and 2006. Based on these trends, fishing effort and harvest opportunity in the directed commercial fishery appeared to peak in 1994-1995, and again in 2005-2006.




Figure 1. Trends in fishing opportunity, drift fishing effort, and king salmon harvest in the directed commercial fishery, 1992-2021.

From 1992-2010, annual commercial harvests ranged from just over 11,000 (1999) to nearly 119,000 (1994) king salmon and exhibited a general declining trend (Figure 2). Directed fishery harvests during this period varied greatly, comprising from 3% (2008) to 98% (1994) of the total commercial harvest during any given year (average 48%). Directed fishery harvests 1992-1998 comprised a much greater proportion (77% average) of the seasonal harvest than any other period since except for 2002 (85%). From 2003-2006 the directed fishery comprised 43% of the seasonal harvest - still much higher than the 5% average experienced 2007-2010. Across all years since 1992 during which a directed fishery occurred, harvests in the directed fishery comprised an average of 45% of the total season harvest.




Figure 2. Commercial harvests of king salmon in the Nushagak District, 1992-2021.

The directed commercial fishery waned considerably after the 2010 season. The ADF&G ceased issuing a pre-season forecast for king salmon beginning 2011 (Jones et al., 2012). After experiencing a poor run in 2010 and lacking a reliable forecast, managers employed a conservative strategy for the next several years whereby fishing would be scheduled only if a harvestable surplus could be projected using inseason escapement rates. The directed fishery was re-opened in 2013 and 2014 but participation and harvests were relatively low. Indications of a strong run exhibited early in the 2014 season were followed by very poor abundance in the second half and failed to indicate the weak run that ultimately resulted.

Strong sockeye salmon run forecasts for the Nushagak and Wood rivers increasingly factored into management of the Nushagak District beginning in 2015, whereby fishing for sockeye salmon was planned to begin earlier in June to control sockeye salmon escapement (Jones et al., 2016). The directed fishery has not been initiated since 2014 due to poor runs experienced 2010-2014, lack of a pre-season forecast to guide any early season fishing, and the expected increased potential for incidental harvest of king during large sockeye runs.

Incidental harvests of king salmon taken during the commercial fishery for sockeye comprised 55% of the annual king salmon commercial fishery harvest, on average, during years when the directed fishery was opened. During these years, incidental harvests ranged from 5,900 to 72,200 and averaged 22,700 king salmon (Figure 3). During years when the directed fishery was not opened, 4,100 to 49,000 king salmon (average 21,600) were harvested incidentally. From 1992 to 2002, the annual incidental harvest averaged 13,800 and ranged from 5,900 to 25,300 king salmon. Since 2003, the annual incidental harvest in the commercial sockeye fishery averaged 27,200 and ranged from 4,100 to 49,300. The higher incidental king salmon catches in the latter period are likely due to a combination of factors, including a shift from king salmon that would have historically been caught in directed fishing effort to occurring in the sockeye fishery, generally larger sockeye returns

resulting in earlier and more intensive fishing directed at sockeye salmon, and in a few years, due to very early sockeye runs (e.g., 2003, 2013).

Large sockeye runs (~10 million+) observed since 2014 have contributed to increased king salmon harvest levels. King salmon run size is also a factor. However, care should be taken in characterizing apparent trends in the incidental harvest and total return given the uncertainty that exists in escapement estimates, which comprise a large component of the total run during low run years. Of note, commercial harvests of king salmon during the 2020 and 2021 seasons were the 3rd lowest and lowest reported since the Plan was adopted.




Figure 3. Number of king salmon harvested incidentally during the commercial sockeye season, 1992-2021.

Since the NMCSP was adopted in 1992, sockeye runs to the Wood, Nushagak and Igushik Rivers have increased over time (Figure 4; Table 4). Average run sizes increased from 6.5 million sockeye salmon in the 1990s, to 9.4 million (2000-2010) to 13.1 million (2011-2020). Runs to the Nushagak district set all-time records in 2006, and again in 2017 and 2018. The 2021 run was the third largest on record.




Figure 4. Nushagak District sockeye salmon runs (district catch and escapement to Nushagak, Wood and Igushik Rivers), 1992-2021.

With both large and early sockeye runs, managers tend to open the commercial fishery earlier in June, and in the case of large runs, schedule fishing time more intensively throughout the season to control sockeye harvest and escapement (Jones et al., 2016). Figure 5 depicts dates on which the Nushagak District opened to commercial fishing for sockeye salmon with drift gillnets, dates on which fishing began on an every-tide basis for the season, and dates on which fishing was extended until further notice. All three sets of dates, particularly season opening dates, exhibit a trend toward earlier starts to the sockeye fishery and intensive fishing regimes. This trend suggests a direct correlation to the increasing sockeye salmon run size in the Nushagak District.




Figure 5. Key dates associated with the annual commercial drift net fishery for sockeye, including the season opening date (blue circle), start date for fishing on an every-tide basis (orange triangle), and dates on which fishing was extended until further notice (green square).

Since the Plan was adopted in 1992, commercial fishing effort appears to have increased based on permit registration statistics. Annual permit registration increased from the 1990s, when the average approximated 320 permits, to the 2000s and 2010s when the average approximated 415 permits (Table 3; Figure 6). Peak daily drift permit registrations showed a similar trend.




Figure 6. Average and peak number of commercial drift net permits registered in the Nushagak District, 1992-2021.

Compounding the increase in effort, the peak registration date also appears to have trended earlier over time (Figure 7), consistent with the increasing size of sockeye runs in recent years.




Figure 7. Peak daily drift permit registration dates, 1992-2021.

Sport Fishery

Regulations

Sport fishing regulations pertaining to Nushagak River king salmon – which consist of Bristol Bay-wide regulations, Nushagak River specific regulations, and Plan provisions - have been modified six times since the Plan was adopted (Table 5). Regulations governing the sport fishery for king salmon have generally become increasingly restrictive, conservative, and complex throughout the life of the Plan.

Most changes consisted of gear restrictions, season closures, bag limit reductions, and imposition of annual limits adopted for a combination of conservation (e.g., spawning season closures) and/or social or allocative reasons (guideline harvest of 5,000 fish). One notable relaxation of restrictive regulations is the most recent change made December 2018 that repealed Plan provisions directing the ADF&G to restrict the sport fishery under inriver run projection scenarios between 55,000-95,000 fish.

Emergency orders were issued during 12 seasons to restrict the sport fishery as directed by the Plan (Table 6). Within the past 15 seasons, the king salmon fishery was restricted inseason for conservation purposes during nine. Bag limit reductions, followed by reductions in the annual limit, were the most common restrictions enacted. Fishing was restricted inseason to catch-and-release during four years (1996, 1997, 2010, and 2019) and the season was closed to fishing for king salmon during two (1999 and 2010). During three of the years when the fishery was restricted (1999, 2011, and 2012), subsequent

increases in the projected inriver run led managers to ease restrictions partially or completely.

Effort

Sport fishing effort for king salmon is concentrated in three areas: the lower Nushagak River near the village of Portage Creek, the middle section of the Nushagak River near the village of Ekwok, and the midsection of the Mulchatna River between the Stuyahok and Kaktuli rivers (Dye and Borden, 2018). Between 1992 and 1997, effort in the Ekwok area was highly variable. Since about 1999, the lower river fishery has steadily expanded upriver to Ekwok and the 2 areas are merging into a single fishery. Most effort for king salmon in the Nushagak River drainage is concentrated near Portage Creek; areas near Ekwok and in the Mulchatna River support lower levels.

Figure 8 and Table 7 depict sport fishing effort in the Nushagak River for all salmon and freshwater species. Dye and Borden (2018) reported that angling for king salmon in the middle section of the Mulchatna River seemed to have diminished since bait was prohibited there in 1992. In the mainstem Nushagak River, effort varied from approximately 10,000 to 20,000 angler days until 2020, the first year of the Covid-19 pandemic, when it fell to 3,400 angler days.




Figure 8. Sport fishing effort (angler-days) in the Nushagak River, 1992-2020.

Based on freshwater logbook data from the period 2006-2018, 41 to 65 (average 51) guide businesses and 155-250 (average 213) guides have operated on the Nushagak River (all species) (Figure 9; Table 8). During any given year, the guide industry served approximately 1,400 to 3,100 clients (average 2,505), many of whom fished for king salmon. Business and guide activity were at their highest early during this period. Like trends observed above for angling effort, the number of guides and businesses declined through about 2010-2012 and then increased to a level slightly lower than that observed in 2006-2007. Guided effort (client days) and harvest followed a very similar trend. Reasons for the decline in participation between 2005-2010 are varied. However, national economic downturns experienced during that time likely played a primary role in the dynamics observed in

guided fishing activity. The ADF&G logbook program was discontinued following the 2018 season.




Figure 9. Number of sport fishing businesses and guides (top), client days (middle), and king salmon harvest by clients (bottom) as estimated by the ADF&G Freshwater Logbook program for the Nushagak River, 2006-2018.

Harvests

Sport harvests of king salmon (guided and unguided) in the Nushagak River ranged from approximately 1,950 (2020) to 10,600 (1994) and averaged 6,130 fish (Figure 10; Table 7). Approximately one-third (39%) of the harvest occurs below the sonar. Like trends in sport fishing effort, annual harvests have varied but have remained generally stable. Prior to 2020, early in the Covid pandemic, no less than 3,500 king salmon were harvested in the fishery during any given year since the Plan was adopted.




Figure 10. Sport harvests of king salmon in the Nushagak River, 1992-2020.

Subsistence Fishery

Regulations, Effort, and Harvest

Nelson (1987) noted that, compared to commercial fishing regulations, few restrictions had been imposed on the subsistence fisheries in Bristol Bay. Of the restrictions that had been enacted prior to the mid-1980s, Nelson noted that the 1974 limit on fishing time (3 days/week) and net length (10 fathoms) on the Dillingham beaches from June 16 to July 17 had the most impact on king salmon harvest rates. Relatively few regulatory changes to the Nushagak subsistence fishery have been enacted since the adoption of the Plan, with two notable exceptions. In 2018, the Board repealed the limits to subsistence fishing periods (i.e., weekly 3-day schedule) and allowed subsistence fishing with dip nets near Dillingham.

Participation in the subsistence fishery (for all salmon species), based on the number of permits issued, appears to have increased steadily but incrementally since adoption of the Plan (Halas and Neufeld, 2018). Comparing average figures for 1992-1996 against those for 2017-2021 indicates the number of subsistence salmon permits issued increased by about 22% (Figure 11, Table 9; Note: estimates for 2020 and 2021 are preliminary). Between the same two time periods, the number of king salmon harvested annually declined by over 38%, and the number per permit decreased by about 49%. Annual harvests and harvest rates began a steady decline in 2018, and in 2020 and 2021 were the lowest since the adoption of the Plan. These recent declines correlate with record large sockeye salmon runs

which have contributed to increased subsistence harvests of sockeye salmon. Both small recent king salmon runs and increased harvests of sockeye salmon in the subsistence fishery likely contributed to the recent decrease in king salmon harvest rates in the subsistence fishery.

Trends in the subsistence fishery, apart from recent low king salmon harvests, are not unlike those observed by Nelson over 30 years ago. He stated then: "Since subsistence fishing is considered a priority use of the resource in Alaska, subsistence use can be expected to continue at near record levels of effort. Harvest levels are expected to remain high, and will continue to be somewhat independent of stock abundance..." It is likely the same outlook holds true today, albeit with a question concerning harvest levels in the near future.




Figure 11. Number of subsistence fishing permits issued (top), estimated king salmon harvest (middle), and harvest per permit (bottom) in the Nushagak District, 1992-2021.

Plan Performance

This section will discuss how the fisheries have performed with respect to management objectives within the Plan.

Changes in Escapement Assessment Tool

Before going further, some discussion is needed regarding the inriver assessment of king salmon because two objectives (inriver run goal and biological escapement goal) rely directly on it and significant uncertainties surround the sonar project and its results.

In 1997, aerial surveys of king salmon spawners raised concern over the accuracy of the sonar counts (Brookover et al., 1997). A distribution study on coho salmon that year coupled with low water conditions indicated that a substantial number of king salmon migrated offshore of the effective reach of the sonar and, as a result, the ADF&G committed to assessing offshore distribution of salmon as an integral component of the project in the future.

Beginning 2002, the ADF&G began using dual frequency identification sonar (DIDSON) concurrently with the Bendix acoustic system then in use (Buck et al., 2012). DIDSON is a type of imaging sonar considered to be generally superior to the 1960s technology used for the Bendix equipment². Comparisons over the next few years found that the DIDSON detected a higher number of fish than the Bendix system, particularly in the more distant-from-shore areas that had been ensonified. In 2005, after a few partial-year, partial-river-segment comparisons of counts from each sonar the ADF&G transitioned to using the DIDSON technology to measure the inriver salmon runs at Portage Creek, and discontinued use of the aging and increasingly difficult-to-service Bendix equipment. Conversion factors for king salmon and other species were subsequently calculated from the relationship between DIDSON and Bendix passage and applied to historical Bendix passage estimates. The revised estimates were then used to produce revised total run and brood tables for Nushagak salmon composed of DIDSON or equivalent estimates.

More recently, ADF&G updated the time series for Nushagak River king salmon to include various sources of historical harvest and escapement data and conducted a run reconstruction and stock recruit analysis using the updated time series (ADF&G Nushagak escapement goal memo, July 11, 2019). During the review, it had become apparent to ADF&G that the run reconstruction and analysis were compromised by a lack of year-to-year overlap among the methods used to estimate escapement. Paired Bendix and DIDSON counts for both riverbanks and multiple years were lacking, Bendix estimates did not align well with paired aerial survey data, and aerial survey data did not overlap in time with DIDSON estimates.

Erickson et al. (2018) summed up uncertainties associated with the current sonar program in a report to the Board in December 2018. A 2011–2014 acoustic tagging study estimated that the sonar beam covered less than a third of the Nushagak River channel. “Preliminary results from the 2011–2014 acoustic tagging study estimated the proportion of king salmon

² In addition, the Bendix equipment was becoming more and more difficult to service and maintain. Al Menin, who invented the Bendix sonar, continued to service the Bendix equipment until 2005.

traveling outside the sonar beam range was 47–65% with a mean of 57%. Similarly, a 2014–2016 mark–recapture study estimated the abundance of adult king salmon in the Nushagak River independently from the sonar estimate. Both studies indicated that a substantial number of king salmon are not enumerated by the existing sonar assessment and that the current sonar assessment is an index of abundance. At this time, ADF&G has not quantified the consistency of the sonar index.”

This assessment of Plan performance takes the current inriver abundance estimates, and resulting spawning escapement and total run estimates, at face value (Table 10). This is problematic in that inriver abundance estimates prior to 2013 were revised by Buck et al. (2012). As a result, management performance in achieving an inriver or escapement goal, for example, can not readily be assessed, at least using the revised estimates, for years prior to 2013. The 1997 season provides a good example of the challenges. In 1997, spawning escapement estimated by aerial surveys (82,000) was twice the sonar count, indicating a problem with the sonar. The revised inriver run estimate presented in Buck et al. (2012) is 170,610. Using the original sonar count, the inriver goal of 75,000 at the time was not met. Using the aerial survey count, the inriver goal was met. And using the current estimate the inriver goal was far exceeded.

Figure 12 and Table 10 depict the Nushagak River king salmon total run estimates. Based on these estimates, runs have generally declined since the Plan was adopted. Recent runs (2016-2020) have averaged about 111,000 fish which is about 42% less than the long-term (1992-2020) average. The most recent three runs (2019-2021) are the smallest since the Plan was adopted. The 2020 king salmon run is the smallest on record, followed by the 2019 run. Once harvest estimates become available for the sport fishery, the 2021 run is likely to replace the 2019 run as the second lowest. Harvest among the fisheries has generally followed the same downward trend throughout the period. This includes the recent three years, and particularly 2020 and 2021 for which total harvests were the lowest observed since the plan was adopted.




Figure 12. Nushagak River king salmon total run and harvest (all fisheries combined), 1992-2020.

Plan Objectives:

The department shall manage the commercial and sport fisheries in the Nushagak District to achieve an inriver goal of 95,000 king salmon in the Nushagak River upstream from the department sonar counter.

Inriver run performance can be assessed by a simple comparison of the estimated inriver run as enumerated at the sonar with the inriver run goal. The combination of changes to the inriver run goal and as stated above, the Bendix-DIDSON conversion makes assessment difficult for years prior to 2013. For this reason, only 2013 through the current year is assessed. Since 2013, the estimated inriver run exceeded the inriver run goal four times but fell short five, including the three most recent years (Figure 13). In 2019-2021, estimated total runs were not large enough to provide for the inriver goal even if no king salmon would have been harvested.




Figure 13. Inriver (and total) run estimates compared to the inriver run goal, 2013-2021.

Provide for a biological escapement goal of 55,000 - 120,000 fish.

Since 2013, estimated spawning escapement fell within the goal range (55,000-120,000 spawners) in five years and fell short in three (Figure 14). Although the spawning escapement estimate is not yet available for 2021, it very likely fell short of the lower bound considering harvests that occur upstream of the sonar, where inriver abundance was estimated at 55,222 king salmon. Aerial surveys conducted in 2017, 2019 and 2021 indicated that actual spawning escapement was likely greater than estimated by sonar; surveys conducted in 2020 seemed to corroborate the low (sonar-based) estimate that year (J. Head, ADF&G, personal communication). From a biological standpoint, the Plan appears to be working generally well in ensuring spawning goals are achieved over the long term. However, should future king salmon runs continue near current levels, achieving inriver goals will likely pose a continued challenge.




Figure 14. Spawning escapement (and total run) estimates compared to the escapement goal (55,000-120,000), 2013-2021.

Provide for reasonable opportunity for subsistence harvest of king salmon; and a king salmon sport fishery guideline harvest level of 5,000 fish, 20 inches or greater in length.

King salmon harvests have declined in the commercial fishery and have remained relatively stable in subsistence and sport fisheries until 2020, when harvests in both fisheries sharply declined (Figure 15; Table 10).




Figure 15. Trends in harvests of Nushagak River king salmon among the commercial, subsistence and sport fisheries, 1992-2021.

The sport fishery guideline harvest level (5,000 king salmon) applies when projected inriver runs do not exceed the inriver goal of 95,000 king salmon. Since 2013, inriver run estimates fell at or below the inriver run goal in 5 years: 2014, 2017 and 2019-2021. Sport harvest estimates are not available for 2021. Harvests in the remaining four years exceeded the guideline harvest level in three years (2014, 2017 and 2019) and fell below in 2020.

Maintain a natural representation of age classes in the escapement.

The Plan's objective to maintain a natural representation of age classes in the escapement has not been addressed in this analysis. Nor has the objective of providing reasonable opportunity for subsistence harvest of king salmon. Addressing the first was beyond the time available to prepare this draft report. The second was beyond the scope. Both, however, are core Plan objectives and should be assessed.

Management Challenges

Many of the recommendations Nelson made in 1987 have been partially or fully carried out. A biological escapement goal was developed in 1992 and subsequently refined in 2012. Development of the Portage Creek sonar has continued through conversion to DIDSON technology, which expanded the portion of the river width ensonified, and the commercial fishery is managed as recommended – by emergency order and using mesh size restrictions to reduce catch rates and achieve a better distribution of escapement through time.

However, several challenges Nelson identified in 1987 – inriver run abundance assessment, overlap between king salmon and sockeye salmon run timing, and size selectivity - remain today. More recently, dynamics have emerged creating new types of challenges. Large record-setting sockeye runs to the Wood and Nushagak Rivers have coincided with poor king salmon runs and exacerbated the difficulties inherent to managing the two species for independent inriver abundance goals. Recent tagging studies and aerial surveys cast considerable uncertainty on the use of sonar-based inriver abundance estimates for managing the Nushagak River fisheries and raised questions after-the-fact on some restrictions predicated on the sonar.

To address these challenges and develop comprehensive recommendations to the Board, the working committee met on numerous occasions over the past three years and discussed possible changes to the NMKSMP for consideration at the November 2022 Bristol Bay meeting. Findings of the committee, including a more robust assessment of current challenges associated with Nushagak River king salmon, will be presented in a separate report.

ACKNOWLEDGEMENTS

A number of individuals made critical direct contributions to this report, without which it would not have been possible. I would like to thank the following members of the ADF&G for their insight and review of the content and their response to multiple requests for data: Lee Borden, Robin Dublin, Jason Dye, Jack Erickson, Jordan Head, Bronwyn Jones, Terri Lemons, Matt Nemeth, Gayle Neufeld, Bob Powers, and Tim Sands. I would also like to thank Mike Nelson for his review of an early draft of the report. I thank members of the Nushagak King Salmon Committee for their insight and review of the manuscript: Peter Christopher, Robert Heyano, Bud Hodson, Bob Klontz, Brian Kraft, Nanci Morris-Lyons, Tom O'Connor,

and George Wilson Jr.. Thanks go to Michael Link and Jeff Regnart of the Bristol Bay Science and Research Institute for their insight and guidance in the report development. I acknowledge many other individuals who directly or indirectly contributed to the Nushagak River king salmon fisheries, their management programs, and this report.

Literature Cited

Alaska Board of Fisheries Meeting Information [Internet]. 2003-. Anchorage, AK; Alaska Department of Fish and Game, Boards Support Section (cited March 6, 2022). Available from: <https://www.adfg.alaska.gov/index.cfm?adfg=fisheriesboard.meetinginfo>

Alaska Department of Fish and Game Emergency Orders and Press Releases [Internet]. Anchorage, AK; Alaska Department of Fish and Game, Division of Sport Fish (cited March 6, 2022). Available from: <https://www.adfg.alaska.gov/sf/EONR/index.cfm>

Alaska Sport Fishing Survey database [Internet]. 1996-. Anchorage, AK; Alaska Department of Fish and Game, Division of Sport Fish (cited October 3, 2019). Available from: <http://www.adfg.alaska.gov/sf/sportfishingsurvey/>

ADF&G (Alaska Department of Fish and Game). 1993. Annual management report, 1992, Bristol Bay area. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A93-32, Anchorage.

ADF&G (Alaska Department of Fish and Game). 1994. Annual management report 1993, Bristol Bay area. Alaska Department of Fish and Game, Commercial Fisheries Management and Development Division, Regional Information Report 2A94-02, Anchorage.

ADF&G (Alaska Department of Fish and Game). 1996. Annual management report 1995, Bristol Bay area. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A96-06, Anchorage.

ADF&G (Alaska Department of Fish and Game). 1997. Annual Management Report 1996 Bristol Bay Area. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A97-14.

ADF&G (Alaska Department of Fish and Game). 1997. Annual management report, 1997, Bristol Bay Area. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A98-08, Anchorage.

ADF&G (Alaska Department of Fish and Game). 1999. Alaska Department of Fish and Game Division of Commercial Fisheries Annual Management, 1998, Bristol Bay. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A99-18, Anchorage.

ADF&G (Alaska Department of Fish and Game). 2001. Annual management report 2000, Bristol Bay area. Alaska Department of Fish and Game, Regional Information Report 2A01-10, Anchorage.

ADF&G (Alaska Department of Fish and Game). 2021. 2021 Bristol Bay Salmon Season Summary. Alaska Department of Fish and Game, Advisory Announcement, September 29, 2021, Anchorage.

Brookover, T.E., R. E. Minard, and B.A. Cross. 1997. Overview of the Nushagak Chinook salmon fishery with emphasis on the sport fishery. Report to the Alaska Board of Fisheries. Regional Information Report No. 2A97-35. Alaska Department of Fish and Game, Anchorage, Alaska.

Browning, J., and J. D. Miller. 1995. Bristol Bay area annual management report, 1994. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A95-11 Anchorage.

Buck, G. B., C. B. Brazil, F. West, L. F. Fair, X. Zhang, and S. L. Maxwell. 2012. Stock assessment of Chinook, sockeye, and chum salmon in the Nushagak River. Alaska Department of Fish and Game, Fishery Manuscript Series No. 12-05, Anchorage.

Dye, J. E., and L. K. Borden. 2018. Sport fisheries in the Bristol Bay Management Area, 2016-2018. Alaska Department of Fish and Game, Fishery Management Report No. 18-27, Anchorage.

Elison, T., P. Salomone, T. Sands, M. Jones, C. Brazil, G. Buck, F. West, T. Krieg, and T. Lemons. 2015. 2014 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 15-24, Anchorage.

Elison, T., P. Salomone, T. Sands, G. Buck, K. Sechrist, and D. Koster. 2018. 2017 Bristol Bay annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 18-11, Anchorage.

Erickson, J. W., G. B. Buck, T. R. McKinley X. Zhang, T. Hamazaki, and A.B. St. Saviour. 2018. Review of salmon escapement goals in Bristol Bay, Alaska, 2018. Alaska Department of Fish and Game, Fishery Manuscript No. 18-06, Anchorage.

Fair, L. F. 2002. Annual management report, 2001, Bristol Bay area. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A02-18, Anchorage.

Fair, L., D. Crawford, F. West, and L. McKinley. 2004. Annual management report, 2003, Bristol Bay area. Alaska Department of Fish and Game, Regional Information Report 2A04-16, Anchorage.

Halas, G. and G. Neufeld. 2018. An Overview of the Subsistence Fisheries of the Bristol Bay Area. Alaska Department of Fish and Game Division of Subsistence, Special Publication No. BOF 2018-04, Anchorage.

Howe, Allen L., Gary Fidler, Allen E. Bingham, and Michael J. Mills. 1996. Harvest, catch, and participation in Alaska sport fisheries during 1995. Alaska Department of Fish and Game, Fishery Data Series No. 96-32, Anchorage.

- Howe, Allen L., Gary Fidler, and Michael J. Mills. 1995. Harvest, catch, and participation in Alaska sport fisheries during 1994. Alaska Department of Fish and Game, Fishery Data Series No. 95-24, Anchorage.
- Jones, M., T. Sands, S. Morstad, P. Salomone, T. Baker, G. Buck, and F. West. 2009. 2008 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 09-30, Anchorage.
- Jones, M., T. Sands, S. Morstad, T. Baker, G. Buck, F. West, P. Salomone and T. Krieg. 2012. 2011 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 12-21, Anchorage.
- Jones, M., T. Sands, S. Morstad, C. Brazil, G. Buck, F. West, P. Salomone, and T. Krieg. 2013. 2012 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No.13-20, Anchorage.
- Jones, M., T. Sands, C. Brazil, G. Buck, F. West, P. Salomone, S. Morstad, and T. Krieg. 2014. 2013 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 14-23, Anchorage.
- Jones, M., T. Sands, T. Elison, P. Salomone, C. Brazil, G. Buck, F. West, T. Krieg, and T. Lemons. 2016. 2015 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 16-13, Anchorage.
- Mills, M. J. 1994. Harvest, catch, and participation in Alaska sport fisheries during 1993. Alaska Department of Fish and Game, Fishery Data Series No. 94-28, Anchorage.
- Mills, M. J. 1993. Harvest, catch and participation in Alaska sport fisheries during 1992. Alaska Department of Fish and Game, Fishery Data Series No. 93-42, Anchorage.
- Minard, M., J. Skrade, T. Brookover, D. Dunaway, B. Cross, and J. Schichnes. 1992. Escapement requirements and fishery descriptions for Nushagak drainage Chinook salmon. Report to the Alaska Board of Fisheries. Regional Information Report No. 1D91-09. Alaska Department of Fish and Game, Dillingham, Alaska.
- Morstad, S. 2000. Annual management report, 1999. Bristol Bay area. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A00-20, Anchorage.
- Morstad, S., M. Jones, T. Sands, P. Salomone, T. Baker, G. Buck, and F. West. 2010. 2009 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 10-25, Anchorage.
- Nelson, M. L. 1987. History and management of the Nushagak Chinook salmon fishery. Bristol Bay Data Report No. 87-1. Alaska Department of Fish and Game, Dillingham, Alaska.

Salomone, P., S. Morstad, T. Sands, C. Westing, T. Baker, F. West, and C. Brazil. 2007. 2006 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 07-22, Anchorage.

Salomone, P., S. Morstad, T. Sands, M. Jones, T. Baker, G. Buck, F. West, and T. Kreig. 2011. 2010 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 11-23, Anchorage.

Salomone P., T. Elison, T. Sands, G. Buck, T. Lemons, F. West, and T. Krieg. 2017. 2016 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 17-27, Anchorage.

Salomone, P., T. Elison, T. Sands, J. Head, and T. Lemons. 2019. 2018 Bristol Bay annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 19-12, Anchorage.

Sands, T., C. Westing, P. Salomone, S. Morstad, T. Baker, F. West, and C. Brazil. 2008. 2007 Bristol Bay area annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 08-28, Anchorage.

Tiernan, A., T. Elison, T. Sands, J. Head, S. L. Vega, and T. Lemons. 2021. 2019 Bristol Bay annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 21-04, Anchorage.

Tiernan, A., T. Elison, T. Sands, J. Head, S. L. Vega, and G. Neufeld. 2021. 2020 Bristol Bay annual management report. Alaska Department of Fish and Game, Fishery Management Report No. 21-16, Anchorage.

Weiland, K. A., S. Morstad, J. B. Browning, T. Sands, L. Fair, D. Crawford, F. West, and L. McKinley. 2003. Annual management report -2002- Bristol Bay. Alaska Department of Fish and Game, Division of Commercial Fisheries, Regional Information Report 2A03-18, Anchorage.

Westing, C., T. Sands, S. Morstad, P. Salomone, L. Fair, F. West, C. Brazil, and K. A. Weiland. 2006. Annual management report 2005 Bristol Bay area. Alaska Department of Fish and Game, Fishery Management Report No. 06-37, Anchorage.

Westing C., S. Morstad, K. A. Weiland, T. Sands, L. Fair, F. West, and C. Brazil. 2005. Annual Management Report 2004 Bristol Bay Area. Alaska Department of Fish and Game, Fishery Management Report No. 05-41, Anchorage.

Appendix A. 1992 Version, Nushagak-Mulchatna Chinook Salmon Management Plan.

5 AAC 06.361. NUSHAGAK-MULCHATNA CHINOOK SALMON MANAGEMENT PLAN. (a) The purpose of this management plan is to ensure adequate spawning escapement of chinook salmon into the Nushagak-Mulchatna river systems. It is the intent of the Board of Fisheries that Nushagak-Mulchatna chinook salmon be harvested in the fisheries that have historically harvested them. The plan in this section provides management guidelines to the department in an effort to preclude allocation conflicts between the various users of this resource. The department shall manage Nushagak-Mulchatna chinook salmon stocks in a conservative manner consistent with sustained yield principles and the subsistence priority.

(b) The department shall manage the commercial fishery in the Nushagak District to achieve an inriver goal of 75,000 chinook salmon present in the Nushagak River upstream from the department sonar. The inriver goal provides for:

- (1) a biological escapement requirement of 65,000 fish;
- (2) reasonable opportunity for subsistence harvest, and;
- (3) a chinook salmon sport fishery harvest of not more than 5,000 fish.

(c) If the total inriver chinook salmon return in the Nushagak River is projected between 75,000 and 95,000 fish, the inriver chinook salmon sport fishery harvest shall not exceed 6,000 fish.

(d) If the total inriver chinook salmon return in the Nushagak River is projected to be between 40,000 and 74,999 fish, the department shall;

(1) by emergency order, close the directed chinook salmon commercial fishery in the Nushagak District; during a closure under this paragraph, the use of a commercial gillnets with webbing larger than 5 1/2 inches, is prohibited; and

(2) if the projected inriver return of chinook salmon in the Nushagak River is less than 65,000 fish, restrict the chinook salmon sport fishery in the Nushagak River by establishing periods by emergency order during which, at the departments discretion, one or more of the following is in effect;

- (A) bag and possession limits are reduced to one (1) fish;
- (B) the use of bait is prohibited;
- (C) time or area for fishing is reduced;
- (D) the chinook salmon sport fishery is closed.

(e) If the total inriver chinook salmon return in the Nushagak River is projected to be less than 40,000, the department shall;

(1) close the sockeye salmon commercial fishery in the Nushagak District until the projected sockeye salmon escapement into the Wood River exceeds 100,000 fish;

(2) close the sport fishery in the Nushagak River to the taking of chinook salmon; and

(3) by emergency order, establish periods during which time or area is reduced for the inriver chinook salmon subsistence fishery in the Nushagak River.

Appendix B. 2019 Version, Nushagak-Mulchatna King Salmon Management Plan.

5 AAC 06.361. Nushagak-Mulchatna King Salmon Management Plan (a) The purpose of this management plan is to ensure biological spawning escapement requirements of king salmon into the Nushagak-Mulchatna river systems. It is the intent of the Alaska Board of Fisheries (board) that Nushagak-Mulchatna king salmon be harvested in the fisheries that have historically harvested them. This management plan provides guidelines to the department in an effort to preclude allocation conflicts between the various users of this resource. The department shall manage Nushagak-Mulchatna king salmon stocks in a conservative manner consistent with sustained yield principles and the subsistence priority.

(b) The department shall manage the commercial and sport fisheries in the Nushagak District as follows:

(1) to achieve an inriver goal of 95,000 king salmon present in the Nushagak River upstream from the department sonar counter; the inriver goal provides for

(A) a biological escapement goal of 55,000 - 120,000 fish;

(B) reasonable opportunity for subsistence harvest of king salmon; and

(C) a king salmon sport fishery guideline harvest level of 5,000 fish, 20 inches or greater in length;

(2) in order to maintain a natural representation of age classes in the escapement, the department shall attempt to schedule commercial openings to provide pulses of fish into the river that have not been subject to harvest by commercial gear;

(3) the department may close the commercial drift or set gillnet fishery if the harvest in the directed commercial king salmon fishery for either gear group is more than two sockeye salmon for every one king salmon.

(c) If the total inriver king salmon return in the Nushagak River is projected to exceed 95,000 fish, the guideline harvest level described in (b)(1)(C) of this section does not apply. (d) If the spawning escapement of king salmon in the Nushagak River is projected to be more than 55,000 fish and the projected inriver return is less than 95,000 fish, the commissioner

(1) shall close, by emergency order, the directed king salmon commercial fishery in the Nushagak District; during a closure under this paragraph, the use of a commercial gillnet with webbing larger than five and one-half inches in another commercial salmon fishery is prohibited;

(2) repealed 5/31/2019;

(3) repealed 5/31/2019;

(e) If the spawning escapement of king salmon in the Nushagak River is projected to be less than 55,000 fish, the commissioner

(1) shall close, by emergency order, the sockeye salmon commercial fishery in the Nushagak District until the projected sockeye salmon escapement into the Wood River exceeds 100,000 fish;

(2) shall close, by emergency order, the sport fishery in the Nushagak River to the taking of salmon and prohibit the use of bait for fishing for all species of fish until the end of the king salmon season specified in 5 AAC 67.020 and 5 AAC 67.022(g); and

(3) shall establish, by emergency order, fishing periods during which the time or area is reduced for the inriver king salmon subsistence fishery in the Nushagak River.

(f) Notwithstanding 5 AAC 06.200, in a directed king salmon commercial fishery, the southern boundary of the Nushagak District is a line from an ADF&G regulatory marker located at Etolin Point at 58° 39.37' N. lat., 158° 19.31' W. long., to 58° 33.92' N. lat., 158° 24.94' W. long. to Protection Point at 58° 29.27' N. lat., 158° 41.78' W. long.

(g) During a directed king salmon commercial fishery in the Nushagak District, drift gillnet and set gillnet fishing periods will be of equal length, but do not have to be open concurrently.

Appendix C. Tables.

Table 1. A chronology of regulatory changes to the Nushagak-Mulchatna River King Salmon Management Plan, 1992-2021.

Year	Modification
1992	Nushagak and Mulchatna King Salmon Management Plan (5 AAC 06.361; Appendix A) is adopted.
1994	Set the sport harvest allocation of 5,000 as a guideline harvest rather than a cap.
1997	Modified the plan directing the department to attempt to schedule commercial openings to provide pulses of chinook salmon into the river that have not been exposed to commercial gear.
	Established an escapement projection of 55,000 king salmon below which inseason restrictions in the sport fishery must be imposed.
2001	Allowed a catch-and-release fishery when the final inriver abundance is projected to be below 55,000 fish but above 40,000 fish. When the king salmon sport fishery is restricted to catch-and-release or is closed for conservation, the use of bait must be prohibited.
2003	Modified provision (d) directing the department to reduce the sport fishing bag limit to 1 per day and in possession, any size, if the projected inriver return falls between 55,000 and 75,000 king salmon.
	Added provision allowing the department to close the commercial drift or set gillnet fishery if the harvest in the directed commercial fishery for either gear group is more than two sockeye salmon for every one king salmon.
2006	Provision added to require, during a directed commercial opening, drift and set gillnet fishing periods to be of equal length, but do not have to be open concurrently.
2012	Modified the biological escapement requirement, inriver goal, and management triggers to reflect changes in inriver sonar operations (Bendix to DIDSON conversion).
2018	Repealed provisions (d)(2) and (3) directing the department to restrict the sport fishery if the projected inriver return falls between 55,000 and 95,000 king salmon.

^a Source: Dye & Borden (2018), Alaska Board of Fisheries Meeting Information [Internet].

Table 2. Fishery statistics for Nushagak District commercial fishing periods targeting king salmon (directed king salmon openings), 1992-2021. All data are preliminary, as reported in annual management reports.

Year	Number of Openings	Opening Duration (Hrs)	Peak Drift Boat Count	# of Deliveries		# Chinook Harvested (Drift & Set)		Source
				Drift	Set	Directed Fishery	Entire Season	
1992	4	32	200			33,905	47,897	ADF&G (1993)
1993	3	23	211			39,536	62,294	ADF&G (1994)
1994	5	122.5	290			111,886	118,643	Browning and Miller (1995)
1995	7	70	347			64,745	80,180	ADF&G (1996)
1996	4	34	252			56,256	73,365	ADF&G (1997)
1997	2	16	278			39,003	64,294	ADF&G (1998)
1998	5	40	-			97,169	108,486	ADF&G (1999)
1999	1	6	279	125	23	563	11,008	Morstad (2000)
2000	-	-		-	-	-	12,055	ADF&G (2001)
2001	-	-		-	-	-	11,050	Fair (2002)
2002	4	30	-	519	594	33,447	39,382	Weiland et al. (2003)
2003	2	11	-	140	48	23,008	42,615	Fair et al. (2004)
2004	2	9	-	153	58	21,233	93,414	Westing et al. (2005)
2005	7	48	-	731	100	30,003	61,854	Westing et al. (2006)
2006	9	66 ^a	-	1,000	194	40,503	83,679	Salomone et al. (2007)
2007	6	74	-	125	2	2,049	51,350	Sands et al. (2008)
2008	2	24	-	26	-	496	18,634	Jones et al. (2009)
2009	3	27	-	122	156	2,575	24,058	Morstad et al. (2010)
2010	3	21	-	33	35	1,143	25,580	Salomone et al. (2011)
2011	-	-		-	-	-	29,811	Jones et al. (2012)
2012	-	-		-	-	-	11,501	Jones et al. (2013)
2013	1	5		8	9	518	15,175	Jones et al. (2014)
2014	4	26 ^b		197	49	3,985	11,448	Elison et al. (2015)
2015	-	-		-	-	-	48,968	Jones et al. (2016)
2016	-	-		-	-	-	23,783	Salomone et al. (2017)
2017	-	-		-	-	-	32,194	Elison et al. (2017)
2018	-	-		-	-	-	35,938	Salomone et al. (2019)
2019	-	-		-	-	-	21,509	Tiernan et al. (2021a)
2020	-	-		-	-	-	6,363	Tiernan et al. (2021b)
2021	-	-		-	-	-	4,103	ADF&G (2021)
^a drift and setnet openings managed separately; drift and setnet hours totaled 66 and 108.								
^b drift and setnet openings managed separately; drift and setnet hours totaled 26 and 8.								

Table 3. Annual drift gill net permit registration statistics, Nushagak District commercial fishery, 1992-2021.

	Average Daily Registration		Peak Daily Registration		Peak Date	Source
	Total Permits ^a	Dual Permits	Total Permits ^a	Dual Permits		
1992	317		360		20-Jun	ADF&G (1993)
1993	250		326		14-Jul	ADF&G (1994)
1994	269		304		23-Jun	Browning and Miller (1995)
1995	225		374		16-Jun	ADF&G (1996)
1996	357		465		11-Jul	ADF&G (1997)
1997	386		499		8-Jul	ADF&G (1998)
1998	404		526		10-Jul	ADF&G (1999)
1999	358		383		30-Jun	Morstad (2000)
2000	402		598		13-Jul	ADF&G (2001)
2001	467		705		1-Jul	Fair (2002)
2002	279		465		2-Jul	Weiland et al. (2003)
2003	407		512		3-Jul	Fair et al. (2004)
2004	362		399		8-Jul	Westing et al. (2005)
2005	527		678		25-Jun	Westing et al. (2006)
2006	564		687		4-Jul	Salomone et al. (2007)
2007	475		741		30-Jun	Sands et al. (2008)
2008	354		470		1-Jul	Jones et al. (2009)
2009	342		431		25-Jun	Morstad et al. (2010)
2010	405		453		1-Jul	Salomone et al. (2011)
2011	424		508		1-Jul	Jones et al. (2012)
2012	282		395		30-Jun	Jones et al. (2013)
2013	313	49	372	60	30-Jun	Jones et al. (2014)
2014	389	65	590	119	27-Jun	Elison et al. (2015)
2015	332	53	474	84	26-Jun	Jones et al. (2016)
2016	364	167	518	244	28-Jun	Salomone et al. (2017)
2017	403	167	636	244	30-Jun	Elison et al. (2017)
2018	803	412	1053	548	27-Jun	Salomone et al. (2019)
2019	603	140	861	207	24-Jun	Tiernan et al. (2021a)
2020	402	84	697	168	26-Jun	Tiernan et al. (2021b)
2021	619	151	855	225	27-Jun	Tim Sands, pers. comm.
^a Total permit sum includes dual boat registrations.						

Table 4. Start dates for initial, intensive, and continuous fishing periods in the commercial fishery for sockeye salmon, and total sockeye run, Nushagak District, 1992-2021. All data are preliminary, as reported in annual management reports (See Table 3 for data sources).

Year	Start Date ^a		Intensive Fishing ^b Start Date		Continuous Fishing ^c Start Date		Sockeye Salmon Total Run	
	Drift	Setnet	Drift	Setnet	Drift	Setnet	Pre-season Forecast	Actual
1992	27-Jun	27-Jun	10-Jul	10-Jul	12-Jul	12-Jul	4,600,000	5,187,351
1993	23-Jun	23-Jun	30-Jun	30-Jun	7-Jul	7-Jul	5,100,000	7,624,224
1994	2-Jul	2-Jul	9-Jul	9-Jul	15-Jul	15-Jul	5,300,000	5,881,064
1995	26-Jun	26-Jun	7-Jul	7-Jul			5,300,000	6,704,568
1996	24-Jun	24-Jun	10-Jul	10-Jul			5,800,000	8,303,614
1997	30-Jun	30-Jun ^d					5,700,000	4,639,699
1998	5-Jul	5-Jul	15-Jul	15-Jul			5,300,000	5,402,866
1999	2-Jul	2-Jul	6-Jul	6-Jul			4,900,000	8,533,542
2000	28-Jun	28-Jun		12-Jul	17-Jul	14-Jul	5,490,000	8,484,050
2001	24-Jun	24-Jun		2-Jul	15-Jul	10-Jul	7,800,000	7,289,194
2002	28-Jun	27-Jun ^d		29-Jun			5,200,000	4,538,394
2003	22-Jun	23-Jun	24-Jun	24-Jun	12-Jul	29-Jun	6,700,000	8,907,474
2004	21-Jun	20-Jun	29-Jun	24-Jun ^e	17-Jul	1-Jul	7,300,000	8,232,466
2005	21-Jun	21-Jun	25-Jun	26-Jun ^e	17-Jul	30-Jun	7,400,000	10,090,869
2006	25-Jun	25-Jun	26-Jun	26-Jun	12-Jul	27-Jun	7,500,000	15,923,444
2007	25-Jun	24-Jun	5-Jul	25-Jun	15-Jul	6-Jul	8,900,000	10,604,183
2008	26-Jun	26-Jun	30-Jun	27-Jun	14-Jul	2-Jul	10,410,000	10,160,079
2009	23-Jun	22-Jun	24-Jun	23-Jun ^f	12-Jul	3-Jul	8,930,000	9,988,322
2010	25-Jun	25-Jun	9-Jul	8-Jul	15-Jul	12-Jul	10,600,000	11,100,363
2011	26-Jun	25-Jun	26-Jun	25-Jun ^e	9-Jul	2-Jul	9,500,000	6,922,015
2012	28-Jun	26-Jun	7-Jul	11-Jul	13-Jul	13-Jul	6,800,000	4,098,632
2013	22-Jun	21-Jun	22-Jun	21-Jun	5-Jul	25-Jun	5,100,000	5,648,859
2014	25-Jun	24-Jun	25-Jun	25-Jun	7-Jul	30-Jun	8,900,000	10,171,331
2015	22-Jun	21-Jun	27-Jun	27-Jun	9-Jul	3-Jul	8,100,000	8,987,563
2016	19-Jun	19-Jun	26-Jun	26-Jun	16-Jul	9-Jul	10,300,000	10,569,247
2017	22-Jun	21-Jun	22-Jun	22-Jun	6-Jul	26-Jun	8,300,000	20,027,749
2018	20-Jun	19-Jun	24-Jun	19-Jun	16-Jul	13-Jul	21,200,000	33,755,636
2019	20-Jun	20-Jun	22-Jun	20-Jun	16-Jul	23-Jun	9,990,000	17,794,604
2020	25-Jun	25-Jun	4-Jul	1-Jul	12-Jul	6-Jul	12,030,000	12,656,061
2021	24-Jun	24-Jun	25-Jun	25-Jun ^e	11-Jul	29-Jun	14,760,000	27,637,560

^a Dates represent the day on which the Nushagak Section opened to commercial fishing for sockeye salmon. From 1992-1998, the entire district including Nushagak Section was opened to both gear types. Beginning in 1998, openings were established for each gear type and section independently.

^b Dates represent the day on which fishing began to occur on an every-tide basis, regardless of number of hours fished per tide.

^c Dates represent the day on which fishing was extended 'until further notice' by EO.

^d After July 5 (in both 1997 and 2002), all fishing occurred in the WRSNA; the district did not re-open.

^e A 1-tide break in fishing occurred for the drift fleet (July 5, 2004; June 30, 2005; July 1, 2011, June 29, 2021).

^f Two breaks in fishing occurred for the drift fleet (June 27 and July 8, 2009).

Table 5. A chronology of significant sport fishing regulation changes for the Nushagak and Mulchatna Rivers, 1990-2021.^a

Effective year	Bay-Wide Sport	Nushagak-Mulchatna Sport	Nushagak-Mulchatna King Salmon Plan
1990		Season established from January 1 to July 25 upstream of and including the Iowithla River.	
1992		Gear restricted to single-hook artificial lures for the portion of the Mulchatna River between the Koktuli and Stuyahok rivers.	
1992			Nushagak and Mulchatna King Salmon Management Plan (5 AAC 06.361) is adopted. Sport harvest capped at 5,000 fish; escapement projection of 65,000 established as trigger for inseason restrictions in the sport fishery.
1994			Sport allocation set as aguideline harvest rather than a cap.
1997	Bay-wide annual harvest limit of 5 king salmon was adopted. Guides prohibited from retaining any species of fish while guiding.	Bag and possession limit reduced to 2 king salmon per day, only 1 over 28 inches. Annual harvest limit of 4 king salmon adopted for the entire Nushagak–Mulchatna drainage. Kokwok River and Nushagak River upstream from its confluence with Harris Creek closed to fishing for king salmon. July 31 spawning season closure adopted for Nushagak River drainage downstream of Iowithla River outlet.	Escapement projection of 55,000 king salmon established as trigger below which inseason restrictions in the sport fishery must be imposed.
2001	Anglers prohibited from removing king salmon from the water if the fish were to be released. Bag and possession limit for king salmon under 20 inches of 10 per day is adopted bay-wide except Nushagak drainage.		Allow a catch-and-release fishery when the final inriver abundance is projected to be below 55,000 fish but above 40,000 fish. Stipulates that when the king salmon sport fishery is restricted to catch-and-release or is closed for conservation, the use of bait must be prohibited.
2003		Bag and possession limit for king salmon under 20 inches of 5 per day is implemented on the Nushagak drainage. King salmon under 20 inches do not count toward the annual limit of 4 and are in addition to the bag limit for king salmon 20 inches or longer.	If inriver projections fall below 75,000, a bag limit of 1 per day, 1 in possession, no size limit, is implemented.
2012		From May 1 to July 31 only 1 single-hook or single-hook lure may be used and the use of bait is allowed UNTIL an angler harvests a daily bag limit of king salmon 20 inches or greater in length, then that angler can only fish with 1 UNBAITED, single-hook or single-hook lure for the remainder of that day.	Plan amended to reflect counts from the new dual frequency identification sonar counter.
2018			Repealed provisions (d)(2) and (3) directing the department to restrict the sport fishery if the projected inriver return falls between 55,000 and 95,000 king salmon.
^a Source: Dye & Borden (2018), Alaska Board of Fisheries Meeting Information [Internet].			

Table 6. Emergency orders issued for the sport and subsistence fisheries under direction of the Nushagak-Mulchatna King Salmon Management Plan, 1992-2021.^a

Year	Effective Date	Sport	Subsistence
1992			
1993			
1994			
1995			
1996	Preseason	Preseason: Bag and possession limit reduced from 3, 2 over 28 inches, to one of any size.	
	9-Jul	Catch and release only for king salmon.	
1997	Preseason	Bag and possession limit reduced from 3, 2 over 28 inches, to one of any size.	
	30-Jun	Catch and release only for king salmon.	
1998			
1999	30-Jun	Seasonal limit reduced from 4 to 2 fish.	
	2-Jul	Fishing for king salmon closed.	
	6-Jul	Season re-opened with seasonal limit of 2 fish.	
	2-Jul		Fishing in the Nushagak River drainage reduced to 3 days per week until August 1.
2000			
2001			
2002			
2003			
2004			
2005			
2006			
2007	7-Jul	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size.	
2008			
2009			
2010	27-Jun	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size.	
	30-Jun	Retention and use of bait prohibited.	
	5-Jul	Fishing for king salmon closed, bait prohibited.	
	6-Jul		Fishing in the Nushagak River drainage reduced to 3 days per week until August 1.
2011	24-Jun	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size. Annual limit reduced from 4 to 2 fish.	
	13-Jul	Annual limit restored to 4 fish.	
2012	28-Jun	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size. Annual limit reduced from 4 to 2 fish.	
	3-Jul	Annual limit restored to 4 fish.	
	7-Jul	Bag and possession limit restored to 2, 1 over 28 inches.	
2013			
2014	7-Jul	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size.	
2015			
2016			
2017	23-Jun	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size. Annual limit reduced from 4 to 2 fish.	
2018			
2019	3-Jul	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size. Annual limit reduced from 4 to 2 fish.	
	10-Jul	Retention and use of bait prohibited.	
2020	10-Jul	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size. Annual limit reduced from 4 to 2 fish.	
2021	27-Jun	Bag and possession limit reduced from 2, 1 over 28 inches, to one of any size. Annual limit reduced from 4 to 2 fish.	

^a Source: Alaska Department of Fish and Game Emergency Orders and Press Releases [Internet], Morstad (2000), Salomone et al. (2011).

Table 7. ADF&G Alaska Sport Fishing Survey summary of angler effort and harvest in the Nushagak River, 1992-2020.

Year	Angler Days ^a	Harvest ^b		
		Below Sonar	Above Sonar	Total
1992	10,031	1,844	2,911	4,755
1993	14,168	2,408	3,492	5,899
1994	15,460	4,436	6,191	10,626
1995	16,410	2,238	2,713	4,951
1996	14,736	2,346	3,045	5,390
1997	10,958	931	2,567	3,497
1998	17,480	1,640	4,188	5,827
1999	15,028	934	3,304	4,237
2000	18,285	1,389	4,628	6,016
2001	18,951	1,600	4,299	5,899
2002	13,396	1,193	2,500	3,693
2003	16,834	2,203	3,752	5,955
2004	18,869	2,567	4,339	6,906
2005	20,050	2,863	5,702	8,565
2006	20,045	3,166	4,307	7,473
2007	18,457	3,581	6,088	9,669
2008	14,936	3,305	3,395	6,700
2009	15,051	2,451	3,903	6,354
2010	9,668	1,659	2,248	3,907
2011	11,329	1,542	3,302	4,844
2012	14,973	1,833	4,098	5,931
2013	16,082	1,971	4,714	6,685
2014	17,576	2,369	3,891	6,260
2015	13,766	2,514	4,720	7,234
2016	17,737	3,053	5,358	8,411
2017	13,299	2,834	2,837	5,671
2018	13,705	3,715	4,477	8,192
2019	10,460	3,768	2,538	6,306
2020	3,427	1,496	454	1,950
Mean 92-96	14,161	2,654	3,670	6,324
Mean 16-20	11,726	2,973	3,133	6,106
Mean 92-20	14,868	2,340	3,792	6,131

^a 1996-2020; Alaska Sport Fishing Survey database [Internet], 1995; Howe et al.(1996), 1994; Howe et al.(1995), 1993; Mills (1994), 1992; Mills (1993). Only estimates for Nushagak River proper were included, i.e. estimates exclude Mulchatna and Nuyakuk Rivers.

^b 1992-2017; Dye and Borden (2018), 2018 and 2019; Jason Dye personal communication, 2020; Lee Borden personal communication.

Table 8. ADF&G Freshwater logbook summary of guided sport fishing in the Nushagak drainage, 2006-2018.

Year	Businesses	Guides	Trips	Clients ^a	Client Days	Crew Days ^b	Harvest ^c
2006	65	247	3,422	2,971	9,960	395	
2007	62	250	3,147	2,891	9,111	124	4,324
2008	60	240	3,140	2,836	9,259	143	4,621
2009	52	183	2,163	1,931	6,309	124	3,030
2010	47	155	1,697	1,401	4,715	136	1,567
2011	47	168	1,864	1,895	4,970	74	2,140
2012	46	189	2,504	2,299	7,105	102	3,827
2013	47	217	2,932	2,553	8,096	174	3,823
2014	51	215	3,066	2,883	8,760	181	4,095
2015	50	227	3,492	3,091	9,903	193	4,613
2016	53	234	3,186	2,770	8,934	159	4,273
2017	48	218	2,468	2,395	6,878	125	2,925
2018	41	223	2,786	2,644	7,827	136	4,647
Mean	51	213	2,759	2,505	7,833	159	3,657
^a Clients excludes youth anglers and anglers without a sport fishing license written. Crew is also excluded, since they aren't clients.							
^b Crew days are the number of days crew fished and excludes client days.							
^c Source: 2006-2016; Dye and Borden (2018), 2017 and 2018; Jason Dye personal communication.							

Table 9. Nushagak Bay watershed subsistence fishery parameter estimates, 1992-2021.^a

Year	Subsistence Permits Issued	King Salmon Harvest	Harvest/ Permit
1992	476	13,588	29
1993	500	17,709	35
1994	523	15,490	30
1995	484	13,701	28
1996	481	15,941	33
1997	538	15,318	28
1998	562	12,258	22
1999	548	10,057	18
2000	541	9,470	18
2001	554	11,760	21
2002	520	11,281	22
2003	527	18,686	35
2004	511	15,610	31
2005	502	12,529	25
2006	461	9,971	22
2007	496	13,330	27
2008	571	12,960	23
2009	530	12,737	24
2010	528	9,150	17
2011	525	12,461	24
2012	517	10,350	20
2013	582	11,567	20
2014	581	16,049	28
2015	591	12,117	21
2016	649	16,576	26
2017	563	11,122	20
2018	589	12,206	21
2019	620	10,206	16
2020	585	8,350	14
2021	656	5,349	8
Mean 92-96	493	15,286	31
Mean 17-21	603	9,447	16
Mean 92-21	544	12,597	23

^a Source: 1992-2015; Halas and Neufeld (2018), 2016-2019; Gayle Neufeld, ADF&G, personal communication, 2020-2021; Terri Lemons, ADF&G, personal communication. Estimates include the Nushagak, Wood, Snake and Igushik River drainages. 2020 and 2021 data is preliminary.

Table 10. King salmon commercial, subsistence, and sport harvest, and escapement for the Nushagak River drainage, 1992-2021.^a

Year	Total Run	Harvests Below Sonar			Inriver Sonar Estimate	Harvests Above Sonar		Spawning Escapement ^b
		Commercial	Subsistence	Sport		Subsistence	Sport	
1992	232,103	47,563	10,322	1,844	172,374	2,498	2,911	166,965
1993	283,393	62,979	14,498	2,408	203,508	2,919	3,492	197,098
1994	334,606	119,480	11,048	4,436	199,643	3,331	6,191	190,121
1995	271,127	79,943	10,800	2,238	178,146	2,419	2,713	173,014
1996	193,141	72,123	10,217	2,346	108,456	3,063	3,045	102,348
1997	247,327	64,390	11,397	931	170,610	2,981	2,567	165,062
1998	371,638	117,820	7,717	1,640	244,461	4,429	4,188	235,845
1999	149,248	11,178	7,450	934	129,686	2,477	3,304	123,906
2000	138,044	12,120	7,247	1,389	117,288	1,979	4,628	110,682
2001	213,306	11,746	7,972	1,600	191,988	3,372	4,299	184,317
2002	229,485	40,039	6,946	1,193	181,307	4,103	2,500	174,704
2003	225,594	43,485	13,399	2,203	166,507	4,448	3,752	158,307
2004	356,240	100,846 ^c	10,644	2,567	242,183	4,422	4,339	233,422
2005	307,701	62,764	7,951	2,863	234,123	4,471	5,702	223,950
2006	218,861	84,881	6,131	3,166	124,683	3,012	4,307	117,364
2007	125,435	51,831	9,564	3,581	60,459	3,411	6,088	50,960
2008	128,752	18,968	9,149	3,305	97,330	2,571	3,395	91,364
2009	117,936	24,693	9,312	2,451	81,480	2,796	3,903	74,781
2010	94,245	26,056	6,345	1,659	60,185	1,845	2,248	56,092
2011	145,232	26,927	8,485	1,542	108,278	2,981	3,302	101,995
2012	195,106	11,952	7,236	1,833	174,085	2,398	4,098	167,589
2013	132,782	10,213	6,889	1,971	113,709	4,201	4,714	104,794
2014	96,639	11,868	11,942	2,369	70,460	3,890	3,891	62,679
2015	160,713	50,675	9,505	2,514	98,019	2,209	4,720	91,090
2016	167,540	24,937	14,182	3,053	125,368	1,933	5,358	118,077
2017	102,083	33,376	8,912	2,834	56,961	1,827	2,837	52,297
2018	148,007	36,626	10,427	3,715	97,239	1,408	4,477	91,354
2019	80,418	22,725	7,162	3,768	46,763	2,967	2,538	41,258
2020	56,705	7,452	4,725	1,496	43,032	2,265	454	40,313
2021		4,820	3,159		55,222	1,297		
Average								
1992-1996	262,874	76,418	11,377	2,654	172,425	2,846	3,670	165,909
2016-2020	110,951	25,023	9,082	2,973	73,873	2,080	3,133	68,660
1992-2020	190,462	44,471	9,227	2,340	134,425	2,987	3,792	127,647
Percent								
1992-1996		79%	12%	3%		3%	4%	
2015-2019		59%	21%	7%		5%	7%	
1992-2019		71%	15%	4%		5%	6%	

^a Source: 1992-2011 Buck et. al 2012 with the following exceptions: Commercial Harvest data source; ADF&G Fish Ticket Data, Subsistence Harvest data for 2004, 2006, 2008, 2010 and 2011; Jordan Head (ADF&G) personal communication, 2012-2021; Jordan Head (ADF&G) personal communication, 2021 Subsistence Harvest data; Terri Lemons (ADF&G) personal communication.

^b Spawning escapement estimated from inriver sonar abundance less upriver harvest for all years except 1997. 1997 estimate based on aerial surveys that have been expanded to DIDSON Equivalents (Buck et al. 2012).

^c Commercial Harvest includes harvest of 4,087 Chinook salmon that were caught in General District 320-05 as they are most likely of Nushagak origin. (Buck et al 2012)